

Review Article

Review of Anesthesia for ENT, Oral and Maxillofacial Surgery at the National Hospital of Niamey: Situational Analysis in Limited Resource Setting

Boukari BM^{1*}, Maikassoua M², Rabiou MB³, Magai A², Abdoulaye MB⁴, Adakal O⁵, Niandou MA⁶, Moussa MR⁶

¹Department of Anesthesia and Resuscitation at the National Hospital of Niamey-Niger

²Department of Anesthesia-intensive care, Regional Hospital of Maradi, Niger

³Department of Anesthesia and ICU Abubakar Tafawa Balewa University Teaching Hospital Bauchi, Nigeria

⁴Department of Anesthesia-intensive care Regional Hospital of Zinder, Niger

⁵Department of Surgery and surgical specialties, Maradi referral hospital, Niger

⁶Department of Resuscitation anesthesia, Niamey-Niger national hospital, Niger

***Corresponding Author:** Boukari Bawa Mahamane, Faculty of Health Sciences, Abdou Moumouni University of Niamey, Department of Anesthesia and Resuscitation of the National Hospital of Niamey, Niger

Received: 03 November 2021; **Accepted:** 19 November 2021; **Published:** 08 February 2022

Citation: Boukari BM, Maikassoua M, Rabiou MB, Magai A, Abdoulaye MB, Adakal O, Niandou MA, Moussa MR. Review of Anesthesia for ENT, Oral and Maxillofacial Surgery at the National Hospital of Niamey: Situational Analysis in Limited Resource Setting. Anesthesia and Critical Care 4 (2022): 23-32.

Abstract

Introduction

Otorhinolaryngology (ENT) and maxillofacial surgery (MFS) have unique anaesthetic challenges, the main anesthetic concern is sharing of airway. The

aim of this work is to determine the demographic and clinical characteristic of patients in the ENT and MF units at the Niamey National Hospital operated under the period of review.

Methodology

This was a prospective cross-sectional and descriptive study running from January 1 to June 30, 2020. Our study included all patients operated on ENT and MFU during the study period. The parameters studied were socio-demographic, the practice of anesthesia, perioperative critical incidence.

Result

During our study, 147 patients were included out of 2082 admitted to the Niamey National Hospital, accounted for 7.06% of total patients admitted during the study period. ENT accounted for 93 patients and CMF 54. The mean age of our patients was 22 ± 10 years with extremes of 11 months and 85 years. The male to female ratio was 2:1. Surgery was urgent in 14.29% of cases and 46.26% of the patients were referred. Tonsillectomy was the most common indication for ENT procedure. Difficult intubation criteria were detected in 34 patients. The ASA 1 class was in the majority (70.07%). The mean wait time for the entire study population was 5 weeks and 5 days with extremes of 0 days and 486 days. Premedication was indicated among 38 patients, diazepam was the drug administered. All patients had received Rocephine as antibiotic prophylaxis. General Anaesthesia with Endotracheal intubation was the anesthetic technique among 84 patients 90.03% for ENT and 48 patients accounted for 88.8%. Propofol, thiopental, ketamine, and halothane were used as induction and maintenance of anaesthesia. Fentanyl was the only opioid drug. All our patients were extubated on a table with an average duration of anesthesia of 106 minutes with extremes of 20 minutes and 600 minutes. The incident reported

intraoperatively was 11 cases of bleeding. The postoperative period was marked by the occurrence of bronchospasm during extubation in 11 patients and arterial hypotension in 3 others, no mortality was been recorded. In 97.96% of cases, the anesthetic team was made up of Senior Technicians in Anesthesia and Resuscitation, Physician Anaesthetist under the supervision of a consultant anesthetist.

Conclusion

This study demonstrated that safe anaesthesia can be administered for ENT and maxillofacial surgeries in limited resource setting and surgical mortality and morbidity can be reduced, through judicious use of available resources, team work and vigilance

Keywords: Anaesthesia; Otorhinolaryngology; Maxillofacial

1. Introduction

Otorhinolaryngology (ENT) and maxillofacial surgery (MFS) posed a lot of challenges to anesthetists, the main anesthetic concerns includes management of the upper airways due sharing of the airway by both anesthetist and the surgeon, pathology modifying the laryngotracheal structures resulting in difficulties in exposing the glottis amongst others. Proper and thorough evaluation of airway for early detection of difficult airway must be carried out prior to the intervention and a strategy of management established, in order to prevent accidents with rapidly serious consequences [1]. Anesthesia in ENT and CMF poses a wide variety of problems; it interests both adults and children; the acts performed are sometimes simple and very short or long and delicate [2]. In Niger, the Niamey National Hospital is the

highest point in the country's health pyramid. It is the main center where ENT and maxillofacial surgery is performed. The aim of this study was to evaluate the anesthetic techniques for ENT and CMF and outcomes in our institution over the period under review.

2. Methodology

This was a prospective cross-sectional and descriptive study conducted over six months from January 1 to June 30, 2020. Were included in our study all patients operated on ENT and MFS during the study period, in emergency or in planned surgery, regardless of the diagnosis, The informed consent of patients was a prerequisite and their anonymity ensured. Only patients who had been operated but did not give their consent were not included. The variables studied were: Sociodemographic pattern, clinical, Anaesthetic technique, surgical procedures, intraoperative and post-operative complications and outcomes. We collected our data using a pre-established survey form. Detailed preanaesthetic evaluation was conducted and appropriate investigation was done. Airway way assessment was done via thyro-chin distance less than 6 cm in 2 patients, cervical circumference greater than 40 in 2 patients, tumors in 9 patients and a Mallampati score greater than 2 in 5 patients was recorded. In total, a risk factor for difficult intubation assessed core was not achieved in patients less than 10 years old, patients were classified base on ASA classification, Anesthesia: Premedication was indicated among 38 patients using diazepam. All patients had received rocephine as antibiotic prophylaxis. Pre-oxygenation was performed in all patients. The anesthetic

induction was intravenous or inhalational, suxamethonium as muscle relaxant was used among some the patients to facilitated endotracheal intubation while those who supraglottic device had had anaesthesia induced with intravenous agent and maintain on halothane in oxygen. Halothane was the only halogenated hypnotic used for maintenance at 1-1.5% in 100% oxygen, fentanyl was the only opioid analgesic used in all the cases. The parameters monitored intraoperatively were non-invasive blood pressure, heart rate, respiratory rate and oxygen saturation. Urine out was monitored where indicated, blood lost was estimated in all patients and blood was transfused when clinically indicated. All patients were extubated on table following reversal of residual muscle in among the patients paralyzed or LMA removed as appropriate at the end of the procedure on table, patient were handed over to the recovery nurses. Word and Excel software were used as tools for data entry and SPSS version 22.0 was used for analysis.

3. Results

During the study period, we collected data from 147 procedures for ENT and MF surgery out of 2082 surgical procedures performed at the Niamey National Hospital. The overall frequency for ENT and MFS was 7.06%. Among these 147 procedures, 93 were performed in ENT service, frequency of 63.26% while MFS accounted for 54, 36.73%.

The mean age of the patients was 22±10 years with extremes of 11 months and 85 years. The age group of 1 to 5 years constituted about 36 patients accounting for 24.49%. In ENT specialties out of 93

patients, 60 were female, 64.52% with a sex ratio of 2:1. For MFU 54 patients, 28 patients were female 51.85% and a sex ratio of 1:1. More than half of the patients (53.74%) came from the region of Niamey followed by that of Dosso with 14.28%. see table I and II. Table III showed. Our study, 103 patients over 147 were ASA1 (70.07%), among the ENT patients, ENT surgery was the most performed type of surgery (63.26%) with tonsillitis as the main operative indications (55.91%). Table III gives the distribution of patients according to the operative indication and according to the type of surgery.

Sixty nine out of the 94 patients in the study were classified ASA1 74.19%, similarly, for MFS out of 54 patients 34 were classified as ASA1 62.96%. Of the 147 patients in our study a total of 27 33.8% had a medical history. The waiting period In the ENT department the mean waiting period is 35± 28 days, for MFS waiting period interval was 14±11 days. Among the hypnotics used, propofol was more

frequently used 76.3% and 77.8 for ENT and MFS respectively. General Anaesthesia was the anesthetic technique of choice among 84 for ENT and 48 for MFS patients Hypertension accounted for 17.0% of medical history. A history of anesthesia was found in 10.7% and 13% of ENT and MFS patients respectively. In ENT and MFS the duration of surgery were 45±20 minutes and 106±33 minutes respectively. Incidents and accidents during and after surgery: Intraoperative transfusion was necessary in 11 patients in the study population, the postoperative period was marked by the occurrence of bronchospasm during extubation in 11 majority occurs in ENT patients and arterial hypotension in 3 others, no mortality was recorded. All our patients were extubated on a table. The anesthesia team: In 97.96% of cases, the anesthetic team was made up of Senior Technicians in Anesthesia and Resuscitation alone, under the supervision of a resuscitator anesthetist.

Variables	ORL	MFU
Age years	11±18	22±10
Sex		
Male	33(34.5%)	26(48.15%)
Female	60(64.5%)	28 (51.85%)
Occupation		
Employ	42(45.2%)	30(55.6%)
Unemploy	31(33.3%)	13(24.1%)
Student	20(21.5%)	11(20.3%)

Table 1: Distribution of biodata

Origine	ORL	MFS	Percentage
Niamey	60	19	53.74%
Dosso	13	8	14.28%
Tillabéri	0	13	8.04%
Zinder	4	7	7.48%
Tahoua	6	4	6.80%

Maradi	8	0	5.44%
Agadez	2	2	2.72%
Venus d'autres pays	0	1	0.68%
Total	93	54	100%

Table 2: Represents the distribution of patients according to origin

	ORL			CMF	
	Indications Effective operating/type of intervention	%		Indications Effective operating/type of intervention	%
Tonsillitis-VG/ Tonsilectomies	52	55.91	Tumor maxillary/Excision	20	37.04
Goiter/Thyroidectomy	11	11.83	Fracture de Lefort/osteosynthesis	16	29.63
Tumor/resection	9	9.68	Osteitis/sequestrectomy	12	22.22
Nasal Polyp/exérèse	9	9.68	Cyst/resection	11	11.11
Trauma/plasty	6	6.45			
Foreign bodies/extraction	5	5.38			
Laryngeal prothesis installation	1	1.08			
Total	93	100		54	100

Table 3: Distribution of patients according to the surgical indication and type of intervention

Mallampati score	ORL		CMF		Total
	Effective	percentage	Effective	percentage	
1	50	53,76	14	25,93	64
2	14	15,05	19	35,19	33
3	6	6,45	1	1,85	7
4	2	2,15	2	3,70	4
Not assessed	21	22,58	18	33,33	39
Total	93	100	54	100	147

Table 4: Distribution of patients according to the score of Mallampati

Variables	ORL n=93	MFS n=54
Anaesthetic techniques		
GA+ETT	84(90.3%)	48(88.9%)
GA+supraglottic device	4(4.3%)	2(3.7%)
Monitored Anaesth. care	5(5.4%)	4(7.4%)
Induction Agents		

Propofol	71 (76.3%)	42 (77.8%)
Thiopentone	19 (20.4%)	8 (14.8%)
Ketamine	3 (3.2%)	5 (9.3%)
Emergency	11 (11.8%)	8 (14.8%)
Elective	82 (88.1%)	46 (85.2%)
Mean waiting period	35± 28 days	14±11 days
Comorbidity		
Preoperative Hypertension	4 (4.3%)	7 (13%)
Diabetes mellitus	3 (3.2%)	5 (9.2%)
Sickle cell disease	2 (2.1%)	0
Asthma	2 (2.1%)	0
Previous history of Anaesth	10(10.7%)	7 (13%)

Table 5: Anaesthetic technique and Co-rmorbidity

Variables	ORL n=93	MFS n=54
Mean BP mmHg		
Systolic	122	115
Diastolic	54	61
Mean PR per minute	92	88
Mean SPO2 %	98	96
Blood transfusion done	4	7
Adequate urine output	91	49
Intraoperative Incidence		
Hypotension	2 (2.1%)	9 (16.7%)
Hypertension	13 (13.9%)	5 (9.2%)
Tachycardia	10 (10.7%)	18 (33.3%)
Bradycardia	0	0
In advent extubation	0	0
Bronchospasm	0	0
Postoperative Incidence		
Bronchospasm	10(10.7%)	1(1.9%)
Hypotension	1(1.1%)	2(3.8%)
Tachycardia	0	2(3.8%)
Bradycardia	0	0
Shivering	8(8.6%)	11(20.4%)
Unplanned ICU admission	0	0
Duration of surgery	45±20 minutes	106±33 minutes

Table 6: Clinical variables

4. Discussion

Our study focused on the evaluation of anesthetics techniques administered in otorhinolaryngology (ENT) and Maxillofacial surgery (MFS) at the Niamey National Hospital from January 01, 2020 to

June 30, 2020 over a period of 6 months. In this study the overall ENT and CMF anesthetic frequency was 7.06% for total number of surgical anesthesia administered during the study period, with a specific frequency per department of 63.26% for ENT and

36.73% for MFS. This frequency is lower than that of Essola et al. at the Libreville hospital in Gabon in 2013 [3], which had found a frequency of 15.8%. The context in which our study was conducted could justify this difference. In fact, infection with covid-19 had considerably reduced the use of health services by patients. The mean age of patients in our overall population was 22.29 years with extremes of 11 months and 85 years; the age group of 1 to 5 years was the most common at 24.49%. In ENT service the age group of 1 to 5 years was the most represented at 32.26%. In the department of stomatology, it was the age groups of 15 to 30 years and 40 to 65 years that were the most represented with 25.93% each. Cissé B et al. in Mali [4] had a similar average age of 25.82 years but an older age group compared to ours of 15 to 29 years to the tune of 44.1%. On the other hand, this remarkable difference between the majority age groups of our two study sites is easily explained by the types of conditions treated in one or the other specialty. Thus, the ENT department, which has the largest number of our study subjects, generally accommodates childhood pathologies such as adenoids, bronchial foreign bodies, recurrent tonsillitis in children, malformations of the ENT sphere, etc. But also the young age of the Nigerien population, it should be noted that according to a WHO report on health and development in Niger, young people under 15 years old represent more than 50.23% of the general population [5]. While the stomatology department most often accommodates frequent defects in adults. In our series, we noted a predominance of the female sex in ENT department as well as in stomatology with 64.52% and 51.85% respectively and respective sex ratios of 0.55 and 0.92. These results are similar to those found by

Cissé B. at the Nianankoro Fomba hospital in Mali in 2008 [4] and Joachim A. at the Cure hospital in Niamey in 2016 [6], with 66% and 52.50% respectively. Predominantly female, others had found different results: Mamadou B. in his study on anesthetic activities at Kati hospital in 2008 and Bawa B. et al. in Niger in 2001 found a male predominance with respectively 61.6% and 63.35%. [7,8] This predominance of the female sex in our study could be explained by the simple fact that the female sex predominates in the Nigerien population. In our series, hypertension was the most common medical history with 4.76%. Regarding the anesthetic history, 12.24% of our patients had a history of anesthesia, of which 10.20% had general anesthesia; 1.36% spine anesthesia; 0.68% locoregional anesthesia and 87.76% had no known anesthetic history. In ENT department 11.83% had a history of GA and in stomatology 7.41% had a history of GA. These results are similar with several data from the African literature, in particular with those of Mamadou B [7]; who also found a percentage of 17.1 for the medical history and also Moussa BMH. [9] who found 9.9% of a surgical history. On the other hand, our results are lower than those of Abdoulaye D, Coll. in 2006 in Mali [10] which recovered 21.77% of anesthetic antecedent with a predominance of general anesthesia at 33.76%. Study population was classified as ASA1 and 25.85 ASA2. Specifically by department, ENT counted 74.19% of patients classified ASA1 and Concerning the ASA classification at the end of the anesthesia consultation; 70.07% of our in stomatology 62.96% were classified ASA1. These results agree with those of Broulaye C. in Bamako in 2013 [11] in his study which focused on the evaluation of the anesthetic

management of patients operated on under perimedullary anesthesia in the district of Bamako and at the University Hospital of Kati who had found 87.5% of patients classified ASA1 and 12.5% ASA2. On the other hand, they differ from the observations of Moussa B.M.H. to NHN [9], in which the ASA1 class is significantly lower than the ASA2 class at 9.5% and 85.9% respectively. This disparity can be explained by the fact that our study sites (ENT and Stomatology) welcome patients with pathologies that are very rarely life-threatening. The average wait time for our entire population was 5 weeks and 5 days with extremes of 0 days and 486 days. For the ENT service in 63.44% of cases this time was greater than 4 weeks. On the other hand, for the stomatology department, the interval between one and two [1-2] weeks was the most encountered in 33.33% of cases. This observation is far superior to those of many data in African literature, in particular those of Abdoulaye D, Coll. in 2006 in Mali [10] as well as Joachim A. at the Niamey cure hospital in 2016 [6] with respectively 32.04% of cases operated on in less than one week and 65% of patients operated between 7 and 9 days. Our result does not comply with the recommendations of the various anesthesia and resuscitation companies and the WHO, which recommend a duration of at least 72 hours, or 48-72 hours otherwise. The plausible explanation for this is that our study took place while the medical world was reeling from the COVID-19 pandemic coupled with the understaffing mentioned above. As a result, many surgeries had to be postponed for several weeks. In our series for the general population, the most common intervention time interval was 1-3 hours. The average duration of anesthesia was 106 minutes with extremes of 20 minutes and 600

minutes. The same is true for the two specialties; the interval of [1-3h] was predominant with respectively 56.99% and 79.63% in ENT and Stomatology. These results are similar to those of Joachim A. at the Niamey Cure Hospital in 2016 [6], who found an average duration of 110mn with extremes of 30mn and 300mn. However our results are superior to those found by Moussa B.M.H. [9], which reports an average anesthesia time of 73.75mn and extremes of 20mn and 275mn. This very pronounced difference between the duration of anesthesia can be explained by the wide variety of types of operations carried out in the two different specialties. The anesthetic act was performed by senior technicians alone in 97.96% of cases. It should be noted that an anesthetist-resuscitator provides supervision. Essola.L et al in their study entitled Pediatric anesthesia in an African environment at the Libreville hospital center in Gabon in 2013 [3], had found different results with 52.6% of cases where the senior technician performed anesthesia alone. These observations highlight the shortage of anesthetists and resuscitators in our health systems. All of our patients were operated on under GA, of which general anesthesia (GA) with orotracheal intubation was the most used method. More specifically for departments, GA plus Orotracheal Intubation was used 88.17% in ENT and 98.15% in Stomatology. Essola.L et al in Gabon [3] and Moussa B.M.H. in Niger [9] found respectively 98.90% and 96.6% of general anesthesia.

In contrast, Joachim A at the Niamey cure hospital in 2016 [6] reported 55.4% GA combined with ALR. Classical induction had been performed at 96.53%, with propofol as the most widely used intravenous

hypnotic at 76.25%, celocurine used as curare at 76.25% and fentanyl as an opioid in 100% of cases. Abdoulaye D and Coll in 2006 in Mali [10], reported results lower than ours with only 0.85% for propofol and 6.3% for fentanyl. These observations reveal the variability between the anesthetics used in different skies. These could have several explanations; among other things the cost of drugs, availability, but also side effects which depend on the patient's medical condition. Intraoperative transfusion was required in 7.59% of patients in our study population. These blood transfusions in 55.56% of cases were indicated by acute anemia. All the patients were extubated on the table and went through the post-intervention monitoring room. Postoperatively, they had all benefited from postoperative analgesia based on paracetamol. We observed 2 adverse events in patients in the postoperative period 1.36%. It was an arterial hypotension and a case of unquantified but spontaneously favorable postoperative pain. In our study the outcome was favorable in 100% of cases, we had no deaths. Dembelé AS et al at the Pasteur clinic in Bamako in 2013 [12] also found 100% favorable progress and a mortality rate of 0.6% postoperatively.

5. Conclusion

Our study demonstrated that, various anesthetic techniques can be safely administered for ENT and CFM surgical procedures with satisfactory outcomes in limited resource setting. Therefore, careful patient assessment, appropriate anesthetic plan, vigilance and communication remain the key for successful conduct of anesthesia, thus results in better surgical outcomes in limited resource setting like ours.

Niamey National Hospital, need to be upgraded through acquisition of modern equipment such as videos laryngoscopes, Airtrach, respirators with capnography sensor and equipment for the practice of ALR and the presence of anesthetists and resuscitators 24 hours a day. The practice of local and regional anesthesia would be of great help in being able to perform outpatient work in this context where outpatient surgery is desirable in order to reduce hospital stays and ensure good availability of hospital beds.

Conflict of interest

The authors declare no conflicts of interest in connection with this manuscript.

Contribution from the authors

All authors participated in the development of this document.

References

1. Amélie L. Anesthesia for ENT and maxillofacial surgery. Treatise on anesthesia and resuscitation (4th edtn), USA.
2. Jean-Louis B. Anesthesia-Resuscitation in otolaryngology and maxillofacialsurgery (1994).
3. Essola L. Pediatric anesthesia in an African environment: experience of a Gabonese adult hospital (2013).
4. CisséBouare. Anesthesiological activities at the Nianankoro Fomba regional hospital (2008).
5. National Institute of Statistics, Niger (2008).

6. Anatovi Joachim Marc Rolin Nasser. Evaluation of the anesthetic activity at Cure Hospital in Niamey. Doctoral thesis in medicine (2016).
7. Mamadou B. Anesthetic activities at kati hospital: 12-month review, Doctoral thesis in medicine. Bamako (2008)
8. Bawa B. Adverse events of anesthesia in pediatric abdominal surgery. About 191 observations collected in the national hospitals of Niamey and Lamordé. *Annals of Abdou Moumouni University* 12 (2001): 178-184.
9. Moussa Boubacar Mohamed Hayatidiny. Anesthesiological management of abdominal emergencies at the Niamey National Hospital (2013): 15-16.
10. Abdoulaye D, Broulaye SM. Serge Anesthesiology activities at the Gabriel Touré Hospital: Review of the year (2004).
11. Broulaye C. Evaluation of the anesthetic management of patients operated on under perimedullary anesthesia in the district of Bamako and at the University Hospital of Kati (2013): 57.
12. Dembélé AS. Anesthetic activities at the Pasteur medico-surgical and intensive care clinic in Bamako. *Rev. Afr. Anesth. Med. Urg.* 18 (2013): 12-13.



This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)