





































- physical exercise beneficial or a risk? *J Optom* 9 (2016): 139-147.
81. Kyari F, Abdull MM, Wormald R, et al. Risk factors for open-angle glaucoma in Nigeria: results from the Nigeria National Blindness and Visual Impairment Survey. *BMC Ophthalmol* 16 (2016): 78.
  82. Aptel F, Bron A, Lachkar Y, et al. Change in Visual Field Progression Following Treatment Escalation in Primary Open-angle Glaucoma. *J Glaucoma* 26 (2017): 875-880.
  83. Sharifipour F, Panahi-bazaz M, Bidar R, Idani A, and Cheraghian B. Age-related variations in corneal biomechanical properties. *Journal of Current Ophthalmology* 28 (2016): 117-122.
  84. Weinreb R, Robinson M, Dibas M, Stamer W. Matrix Metalloproteinases and Glaucoma Treatment J. of Ocular Pharmacology and Therapeutics 36 (2020): 4.
  85. Prata TS, Lima VC, Guedes LM, et al. Association between corneal biomechanical properties and optic nerve head morphology in newly diagnosed glaucoma patients. *Clin Exp Ophthalmol* 40 (2012): 682-688.
  86. Sharpe GP, Danthurebandara VM, Vianna JR, et al. Optic Disc Hemorrhages and Lamellar Disinsertions in Glaucoma. *Ophthalmology* 123 (2016): 1949-1956.
  87. Johnson CS, Mian SI, Moroi S, Epstein D, Izatt J, Afshari NA. Role of corneal elasticity in damping of intraocular pressure. *Investigative Ophthalmology and Visual Science* 48 (2007): 2540-2544.
  88. Memarzadeh F, Ying-Lai M, Azen SP, et al. Associations with intraocular pressure in Latinos: the Los Angeles Latino Eye Study. *Am J Ophthalmol* 146 (2008): 69-76.
  89. Mansoori T, Sivaswamy J, Gamalapati JS, et al. Measurement of radial peripapillary capillary density in the normal human retina using optical coherence tomography angiography. *J Glaucoma* 26 (2017): 241-246.
  90. Park JH, Yoo C, Girard MJA, et al. Peripapillary Vessel Density in Glaucomatous Eyes: Comparison Between Pseudoexfoliation Glaucoma and Primary Open-angle Glaucoma. *J Glaucoma* 27 (2018): 1009-1016.
  91. Mastropasqua R, Agnifili L, Borrelli E, et al. Optical Coherence Tomography Angiography of the Peripapillary Retina in Normal-Tension Glaucoma and Chronic Nonarteritic Anterior Ischemic Optic Neuropathy. *Curr Eye Res* 43 (2018): 778.
  92. Vysniauskiene I, Allemann R, Flammer J, et al. Vasoactive responses of U46619, PGF2alpha, latanoprost, and travoprost in isolated porcine ciliary arteries. *Invest Ophthalmol Vis Sci* 47 (2006): 295-298.
  93. Xu J, Li Y, Song S, et al. Evaluating changes of blood flow in retina, choroid, and outer choroid in rats in response to elevated intraocular pressure by 1300 nm swept-source OCT. *Microvasc Res* 121 (2019): 37-45.
  94. In JH, Lee SY, Cho SH, et al. Peripapillary vessel density reversal after trabeculectomy in glaucoma. *J Ophthalmol* (2018): 8909714.
  95. Shin JW, Sung KR, Uhm KB, et al. Peripapillary microvascular improvement and lamina cribrosa depth reduction after trabeculectomy in primary open angle glaucoma. *Invest Ophthalmol Vis Sci* 58 (2017): 5993-5999.
  96. Hollo G. Influence of large intraocular pressure reduction on peripapillary OCT vessel density in ocular hypertensive and glaucoma eyes. *J Glaucoma* 26 (2017): e7-e10.
  97. Kuryшева NI, Lepeshkina LV. Selective Laser Trabeculoplasty Protects Glaucoma Progression in the Initial Primary Open-Angle Glaucoma and Angle-Closure Glaucoma after Laser Peripheral Iridotomy in the Long Term. *Hindawi BioMed Research International* (2019).
  98. Zhang X, Dastiridou A, Francis B. et al. Comparison of glaucoma progression detection by optical coherence tomography and visual field. *American Journal of Ophthalmology* 184 (2017): 63-74.