

Research Article

## Evaluation of the Neutrophil-Lymphocyte Ratio in the Prediction of Systemic Infection in Normal Newborns in Lubumbashi: Cross-Sectional Study

Amir Yuma N'Simbo Assumani<sup>1</sup>, Aliocha Natuhoyila Nkodila<sup>2, 3\*</sup>, Jean Lambert Ehungu Gini<sup>4</sup>, Gray Kateng A Wakamb<sup>1</sup>, Gauthier Kasansaika Mutoba<sup>1</sup>, Kasim N'simbo Sangwa<sup>1</sup>, Maguy Omoy Ngongo<sup>1</sup>, Stanislas Okitosho Wembonyama<sup>1</sup>, Oscar Numbi Luboya<sup>1</sup>

<sup>1</sup>Department of Pediatrics, University of Lubumbashi, Lubumbashi, Democratic Republic of Congo

<sup>2</sup>Department of Family Medicine and Primary Health Care, Protestant University in Congo, Kinshasa, Democratic Republic of Congo

<sup>3</sup>Department of Biostatistics and Epidemiology, Lomo University of Research, Kinshasa, Democratic Republic of Congo

<sup>4</sup>Department of Pediatrics, University of Kinshasa, Kinshasa, Democratic Republic of Congo

**\*Corresponding Author:** Aliocha Nkodila Natuhoyila, Department of Family Medicine and Primary Health Care, Protestant University in Congo, Kinshasa, Democratic Republic of Congo

**Received:** 01 June 2022; **Accepted:** 18 June 2022; **Published:** 30 June 2022

**Citation:** Amir Yuma N'Simbo Assumani, Aliocha Natuhoyila Nkodila, Jean Lambert Ehungu Gini, Gray Kateng A Wakamb, Gauthier Kasansaika Mutoba, Kasim N'simbo Sangwa, Maguy Omoy Ngongo, Stanislas Okitosho Wembonyama, Oscar Numbi Luboya. Evaluation of the Neutrophil-Lymphocyte Ratio in the Prediction of Systemic Infection in Normal Newborns in Lubumbashi: Cross-Sectional Study. *Journal of Pediatrics, Perinatology and Child Health* 6 (2022): 305-313.

### Abstract

**Background and Purpose:** Despite progress in the surveillance of newborns in hospitals around the world

in general and the country in particular, systemic infection remains a major concern. The purpose of this study to assess the ability of neutrophil-lymphocyte

ratio (NLR) to predict systemic infection in newborns in clinics in Lubumbashi.

**Methods:** Cross-sectional and analytical study that included 430 normal newborns in 25 medical facilities in the city of Lubumbashi, chosen in a simple random manner during the period from November 2015 to December 2017. The clinical and biological characteristics of newborns were studied.

**Results:** out of 430 children who performed a complete blood count, 106 had an  $NLR > 3$ , a frequency of systemic infection of 24.7%. The mean values of neutrophils, lymphocytes, eosinophils, basophils monocytes were significantly higher in patients with an  $NLR > 3$ . Mean RNL values were  $2.5 \pm 1.2$ , it was  $1.9 \pm 0.6$  in children with  $NLR \leq 3$ , and  $4.3 \pm 1.2$  in those with  $RNL > 3$ . The area under the curve (AUC) for NLR, Neutrophil, Lymphocyte and Platelet were 0.887, respectively; 0.738, 0.639 and 0.552. NLR is more sensitive and specific in predicting systemic infection compared to neutrophil, lymphocyte and platelet count.

**Conclusion:** The results show that NLR is an effective indicator in the diagnosis of systemic infection than neutrophil, lymphocyte and platelet count.

**Keywords:** Systemic Infection; NLR; Normal Newborn; Lubumbashi

## 1. Background

Systemic infection is a generalized disease that progresses rapidly and can lead to high mortality [1, 2]. This infection sets in from a progression of various pathogenic microorganisms that enter the bloodstream, reproduce and then release toxins and

metabolites [2]. The toxins and metabolites released induce a systemic inflammatory response [3]. Currently, sepsis and bacteremia are collectively referred to as systemic infection [3]. In recent years, due to the irrational use of drugs (antibiotics, hormones, immunosuppressants, etc.), chemotherapy and organ transplantation, the incidence of systemic infections has increased [2, 4]. From various invasion pathways of pathogenic microorganisms, patients are witnessed a complex development of systemic infections due to their virulence. Depending on the patient's immune defense, the infection can progress rapidly to complication or death [5]. Early assessment of systemic infections is very important for clinicians as it enables rapid decision-making to finally reduce mortality [6]. The level of white blood cells, neutrophils, lymphocytes, CRP and the neutrophil-to-lymphocyte ratio are common indices used to diagnose a systemic infection [6]. In recent years, the role of the Neutrophil-to-Lymphocyte Ratio has been confirmed in the diagnosis of systemic infections, but they have not been widely used due to controversies [7].

During inflammation, bacteria invade the body, release chemokines in vivo, including the release of cytokines, and then stimulate chemotaxis by producing and releasing large numbers of neutrophils in the bone marrow. This results in elevated local and systemic neutrophilia. Throughout the reaction, the patient's immune system function is diminished, a number of T cells are apoptotic, apoptotic cells inhibit a large number of activated T cells, and the proliferative activity of T cells decreases [8-10]. Peripheral lymphocytes decrease, platelets and neutrophils increase. Therefore, the neutrophil-lymphocyte ratio will also increase. This is why the neutrophil-lymphocyte ratio is often used as a

biomarker of systemic inflammatory disease [9-13]. The purpose of this study to assess the ability of neutrophil-lymphocyte ratio (NLR) to predict systemic infection in newborns in clinics in Lubumbashi.

### 1.1 Population and methods

An analytical cross-sectional study was carried out in 25 medical facilities in the city of Lubumbashi, chosen in a simple random manner during the period from November 2015 to December 2017. The study population was made up of all babies born in maternity hospitals in all of the medical training courses selected. The eligibility criteria for the study were, babies born eutrophic at term, without any complications in the facilities selected during the study period and whose parents had consented to participate in the study. Simple probability and random sampling with a sample step of 3 was used to collect the data for this study. The sample size was calculated from Fisher's formula:  $n \geq (Z^2 \times p(1-p)) / d^2$  where  $n$  = Sample size,  $z = 1,96$  (confidence coefficient),  $p$  = previous prevalence,  $d = 0.05$  (margin of error or range of imprecision reflecting the degree of absolute precision desired). Because of the probable non-responding subjects, 10% of the number calculated at the height should be added. We estimated the frequency of systemic infection to be 50% in infants, as prescribed in the literature, in the absence of such a documented prevalence in the country. The sample size thus calculated was  $n \geq (1.96)^2 \times 0.5 \times 0.5 / (0.05)^2 = 384$ . By including the 10% of non-respondents, we had obtained 422 babies to include. Thus, the distribution of the babies retained was made according to the active file of each Medical Training given the quotas of babies born by selected Medical Training. Data were collected using a structured survey form.

Respondents were admitted consecutively by interview of mothers conducted by the principal investigator and her team. Those who had consented were informed of the purpose of the study and the reasons for which they were approached. An information sheet with detailed explanations was read for them. The following definitions were used in this work: systemic infection, newborn with a Neutrophil-to-lymphocyte ratio  $> 3$  if it is  $NLR \leq 3$ , the newborn has no infection. Bacterial infection, newborn baby with a WBC Zscore or neutrophil  $< -2SD$ .

### 1.2 Statistical analyzes

Analyzes were performed on SPSS 21.0. Descriptive statistics consisted of calculating the mean and standard deviation for quantitative data and proportions for categorical data. Pearson's chi-square test or Fisher's exact test was used to compare the proportions, on the other hand Student's t test compared the means. The prediction of RNL in systemic infection was made using the ROC curves and calculating the area under the curve (AUC). The p value  $< 0.05$  was considered to be the threshold of statistical significance.

### 1.3 Ethical considerations

The study respected the rules of anonymity, confidentiality, justice and the charity of newborns. Beforehand, the protocol for this study was submitted and approved by the Lubumbashi ethics committee under Approval No. UNILU/CEM/070/2017.

## 3. Results

Of 430 children who performed a complete blood count, 106 had an  $NLR > 3$ , for a frequency of systemic infection of 24.7%.

**3.1 General characteristics of the study population**

The sex ratio of the children used was 1H / 1F, no gender predominance. The mean values of neutrophils, lymphocytes, eosinophils, basophils monocytes were significantly higher in patients with an NLR > 3. Mean NLR values were  $2.5 \pm 1.2$ , it was  $1.9 \pm 0.6$  in children with  $NLR \leq 3$ , and  $4.3 \pm 1.2$  in those with  $NLR > 3$ . For the other variables, the differences were not statistically significant (Table 1).

**3.1.1 Prediction of NLR in systemic infection:**


The ROC curves of NLR, Neutrophil, Lymphocyte and Platelet in the prediction of systemic infection are by Figures 1 and 2 and the values interpreting the curves are summarized in Table 2. The area under the curve (AUC) for the NLR, Neutrophil, Lymphocyte and Platelet were 0.887, respectively; 0.738, 0.639 and 0.552. NLR is more sensitive and specific in predicting systemic infection compared to neutrophil, lymphocyte and platelet count.

Variable	Over all n=430	NLR ≤3 n=324	NLR >3 n=106	p
Gender				0.437
Male	228 (53.0)	173 (53.4)	55 (51.9)	
Female	202 (47.0)	151 (46.6)	51 (48.1)	
Birth weight (g)	$3167.1 \pm 430.6$	$3165.4 \pm 436.2$	$3172.1 \pm 414.9$	0.891
Hight (cm)	$48.6 \pm 2.6$	$48.6 \pm 2.6$	$48.8 \pm 2.4$	0.603
Cranial perimeter (cm)	$34.8 \pm 1.4$	$34.8 \pm 1.4$	$34.7 \pm 1.4$	0.439
Thoracic perimeter (cm)	$33.5 \pm 1.4$	$33.5 \pm 1.3$	$33.5 \pm 1.7$	0.941
Gestational age (SA)	$38.6 \pm 1.6$	$38.6 \pm 1.6$	$38.4 \pm 1.7$	0.366
White globule 10 <sup>9</sup> LH	$17.7 \pm 5.4$	$17.0 \pm 5.1$	$19.6 \pm 5.9$	<0.001
Neutrophil 10 <sup>9</sup> LH	$11.1 \pm 4.2$	$9.9 \pm 3.4$	$14.5 \pm 4.5$	<0.001
Lymphocyte 10 <sup>9</sup> LH	$4.9 \pm 1.8$	$5.4 \pm 1.8$	$3.5 \pm 1.1$	<0.001
Eosinophil	$530.1 \pm 162.9$	$510.8 \pm 153.8$	$588.9 \pm 176.1$	<0.001
Basophil	$132.5 \pm 40.7$	$127.7 \pm 38.4$	$147.2 \pm 44.0$	<0.001
Monocyte	$883.5 \pm 271.5$	$851.4 \pm 256.3$	$981.6 \pm 293.6$	<0.001
RNL	$2.5 \pm 1.2$	$1.9 \pm 0.6$	$4.3 \pm 1.2$	<0.001
HB, g/dl	$17.9 \pm 2.4$	$18.0 \pm 2.3$	$17.7 \pm 2.6$	0.260
RBC 10 <sup>12</sup> LH	$6.1 \pm 3.3$	$6.1 \pm 3.8$	$5.8 \pm 0.9$	0.367
TCMH, pg	$30.9 \pm 2.8$	$31.0 \pm 2.9$	$30.8 \pm 2.7$	0.398
CCMH, g/dl	$31.9 \pm 1.8$	$31.8 \pm 1.8$	$32.1 \pm 1.6$	0.114
VGM, fl	$96.6 \pm 8.7$	$96.9 \pm 9.1$	$95.7 \pm 7.2$	0.220
PLT 10 <sup>9</sup> LH	$237.1 \pm 88.1$	$233.2 \pm 85.3$	$248.8 \pm 95.7$	0.114


**Table 1:** General characteristics of the study population according to the category of NLR.

Marqueurs	AUC	SD	p	95% CI	Se	Sp
NLR	0.887	0.036	<0.001	0.816-0.959	0.976	0.898
Neutrophil ( $\times 10^9/l$ )	0.738	0.039	<0.001	0.662-0.814	0.982	0.885
Lymphocyte ( $\times 10^9/l$ )	0.639	0.038	0.001	0.565-0.713	0.891	0.781
Platelet ( $\times 10^6/l$ )	0.552	0.041	0.216	0.470-0.633	0.565	0.491

**Table 2:** Prediction of biomarkers in the diagnosis of systemic infection.



**Figure 1:** ROC curves predicting systemic RNL and neutrophil infection.



**Figure 2:** ROC curves for predicting systemic lymphocyte and platelet infection.

#### 4. Discussion

Systemic infection is a medical emergency requiring early diagnosis and prompt management for the clinician. Delayed treatment increases the mortality rate [10, 14]. We conducted a cross-sectional study in 430 apparently healthy newborns in 24 hospitals in the city of Lubumbashi. The objective of this study was to assess the diagnostic performance of NLR in predicting systemic infection. The salient results showed that NLR better predicted systemic infection, relative to neutrophil, lymphocyte and platelet count.

The mean NLR values of patients with systemic infection were  $4.3 \pm 1.2$ , which is significantly higher than patients without systemic infection ( $1.9 \pm 0.6$ ). Our data are consistent with observations in the literature, including the work of Terradas et al. [15] and Zahorec [8]. It is generally believed that  $NLR > 3.0$  could symbolize positive blood culture results. The critical value of NLR for the diagnosis of systemic infection was set at 3.0, its sensitivity 97.6% and its specificity 89.8%. Additionally, NLR is a good indicator of systemic infection with a subsurface curve of 0.885 (95% CI: 0.816-0.959). It is therefore necessary in poorly equipped settings where blood culture is not feasible, to be satisfied with NLR to enable a systemic infection to be diagnosed earlier.

Neutrophil count predicted systemic infection less, but more better than lymphocyte and platelet count. The sensitivity and specificity of neurophil count in predicting systemic infection was 98.2% and 88.5% and its area under the ROC curve was 0.738 (95% CI: 0.662-0.814). Our results relate to the literature, including the results of Charles et al [16], Brodska et al. [17] and Clec'h et al. [18], but unlike those of Dandona et al. [19], who believes that there is no

significant difference in the prediction of systemic infection between NLR and neutrophil count. For Dandona et al. [19], RNL is more related to the neutrophil level. An elevation in neutrophils should be linearly related to an elevation in NLR because they are closely related. This elevation of neutrophils often observed in systemic infection is due to chemotaxis due to the presence of bacteria in the body during the release of endotoxin as a pro-inflammatory cytokine [20]. On the other hand, Surbatovic et al. had neutrophil and lymphocyte count predicted systemic infection better compared to NLR [21]. The observed difference could be due to the divergence in methods and sample size between the two studies.

There were some limitations to this research. First of all, this is a cross-sectional study, the outcome of which was affected by other factors. We cannot know if other primary diseases, such as neutropenia, would influence the NLR value. Second, the rapid change in the condition of the newborn after birth can cause the NLR to change rapidly, which also affects the accuracy of statistical data.

#### 5. Conclusion

The results of this study show that NLR is an effective indicator in the diagnosis of systemic infection. It could be used as a tool in systemic infection in settings where blood culture is difficult to obtain because its sensitivity and specificity is greater.

#### Declaration

##### Ethics approval and consent to participate

The study protocol was approved by the ethics committee of the University of Lubumbashi (UNILU/CEM/070/2017). The study has been performed in accordance with the ethical standards

laid down in the 1964 Declaration of Helsinki and its later amendments. Written informed consent was requested from the parents or guardian at the beginning of the study. The informed consents included that data from the participants can be published in scientific journals in an anonymized format in which individual responses never are made visible or possible to identify.

### **Consent for publication**

This is not an experimental study but a transversal study and hence, no separate consent for publication is needed.

### **Availability of data and materials**

The data the study is based on contains personal and intimate information that according to initial agreements with the participants from the start of the study cannot be transferred to third parties or outside the borders of Lubumbashi. Theoretically some subsets of data could on reasonable request be delivered but unfortunately, in most cases it probably would be evaluated as not possible along with this. The datasets used and analyzed during the current study are available from the corresponding author on reasonable request: [nkodilaaliocha@gmail.com](mailto:nkodilaaliocha@gmail.com)

### **Competing interests**

The authors declare no conflict of interest.

### **Funding**

This research received no external funding.

### **Acknowledgements**

The authors warmly thank all the investigators who participated in the study, the mothers who agreed to take a sample from their child who was not sick, as

well as the hospitals who accepted the collection of field data.

### **Authors' contribution**

Amir Yuma N'simbo Assumani and Aliocha Natuhoyila Nkodila initiated the study; Jean Lambert Ehungu Gini participated in the study design; Amir Yuma N'simbo Assumani, Aliocha Natuhoyila Nkodila, Oscar Numbi Luboya and Stanislas Okitosho Wembonyama wrote the manuscript; Gray Kateng A Wakamb, Gauthier Kasansaika Mutoba supervised data collection; Kasim N'simbo Sangwa, Maguy Omoy Ngongo and Aliocha Natuhoyila Nkodila validated and analyzed the data; Aliocha Natuhoyila Nkodila interpreted the data; all authors made intellectual contributions to the draft manuscript and approved the final manuscript for submission.

### **References**

1. El Haddad H, Chaftari AM, Hachem R, et al. Biomarkers of sepsis and bloodstream infections: the role of procalcitonin and proadrenomedullin with emphasis in patients with cancer. *Clin. Infect. Dis* 67 (2018): 971-977.
2. Emonet S, Schrenzel J. How could rapid bacterial identification improve the management of septic patients? *Expert. Rev. Anti. Infect. Ther* 9 (2011): 707-709.
3. Ministry of Health P. Diagnostic criteria for nosocomial infections (proposed). *Natl Med. J. China* 81 (2001): 314-320.
4. Li DN, He C, Wang YF, et al. Pathogen characteristics of catheter-related bloodstream infection. *Sichuan Da Xue Xue Bao Yi Xue Ban* 46 (2015): 794-797.
5. Buchanan MO, Summerlin-Long SK,

- Dibiase LM, et al. The compliance coach: a bedside observer, auditor, and educator as part of an infection prevention department's team approach for improving central line care and reducing central line-associated bloodstream infection risk. *Am. J. Infect. Control* 47 (2019): 109-111.
6. Grace CJ, Lieberman J, Pierce K, et al. Usefulness of blood culture for hospitalized patients who are receiving antibiotic therapy. *Clin. Infect. Dis* 32 (2001): 1651-1655.
  7. Azevedo JR, Torres OJ, Czczko NG, et al. Procalcitonin as a prognostic biomarker of severe sepsis and septic shock. *Rev. Col. Bras. Cir* 39 (2012): 456-461.
  8. Zahorec R. Ratio of neutrophil to lymphocyte counts – rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl. Lek. Listy* 102 (2001): 5-14.
  9. Zhao Z, Liu J, Wang J, et al. Platelet-to-lymphocyte ratio (PLR) and neutrophil-to-lymphocyte ratio (NLR) are associated with chronic hepatitis B virus (HBV) infection. *Int. Immunopharmacol* 51 (2017): 1-8.
  10. Yang M, Li L, Su N, et al. Dynamic monitoring of the neutrophil/lymphocyte ratio could predict the prognosis of patients with bloodstream infection. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 27 (2015): 471-476.
  11. Pan YP, Fang YP, Xu YH, et al. The diagnostic value of procalcitonin versus other biomarkers in prediction of bloodstream infection. *Clin. Lab* 63 (2017): 277-285.
  12. Guo D, Jia C, Su H. Eosinophil could predict the prognosis of patients with bloodstream infection: a retrospective analysis of 305 cases. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 29 (2017): 496-500.
  13. Durant SE, Hopkins WA, Davis AK, et al. Evidence of ectoparasite-induced endocrine disruption in an imperiled giant salamander, the eastern hellbender (*Cryptobranchus alleganiensis*). *J. Exp. Biol* 218 (2015): 229.
  14. Ljungstrom L, Pernestig AK, Jacobsson G, et al. Diagnostic accuracy of procalcitonin, neutrophil–lymphocyte count ratio, C-reactive protein, and lactate in patients with suspected bacterial sepsis. *PLoS ONE* 12 (2017): e0181704.
  15. Terradas R, Grau S, Blanch J, et al. Eosinophil count and neutrophil–lymphocyte count ratio as prognostic markers in patients with bacteremia: a retrospective cohort study. *PLoS ONE* 7 (2012): e42860.
  16. Charles PE, Ladoire S, Aho S, et al. Serum procalcitonin elevation in critically ill patients at the onset of bacteremia caused by either Gram negative or Gram positive bacteria. *BMC Infect. Dis* 8 (2008): 38.
  17. Brodska H, Malickova K, Adamkova V, et al. Significantly higher procalcitonin levels could differentiate Gram-negative sepsis from Gram-positive and fungal sepsis. *Clin. Exp. Med* 13 (2013): 165-170.
  18. Clec'h C, Ferriere F, Karoubi P, et al. Diagnostic and prognostic value of procalcitonin in patients with septic shock. *Crit Care Med* 32 (2004): 1166-1169.
  19. Dandona P, Nix D, Wilson MF, et al. Procalcitonin increase after endotoxin injection in normal subjects. *J. Clin. Endocrinol. Metab* 79 (1994): 1605-1608.
  20. Leli C, Ferranti M, Moretti A, et al.

Procalcitonin levels in Gram-positive, Gram-negative, and fungal bloodstream infections. *Dis. Markers* (2015): 701480.

21. Djordjevic D, Rondovic G, Surbatovic M, et al. Neutrophil-to-lymphocyte ratio, monocyte-to-lymphocyte ratio, platelet-to-

lymphocyte ratio, and mean platelet volume-to-platelet count ratio as biomarkers in critically ill and injured patients: which ratio to choose to predict outcome and nature of bacteremia?. *Mediators Inflamm* (2018): 3758068.



This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)